
Syphilis in Escambia County

Syphilis Elimination Task Force Meeting February 12, 2013




2013 Syphilis Cases by Race

African Americans are adversely affected.

- 38 males
- 27 females

White

- 21 males
- 14 females




2012 & 2013 Syphilis Cases by Gender and Race

African American females saw an increase in cases.


African American males saw a decrease, but are still reporting the most cases.

White females saw no change in number of cases.


White males saw a slight decrease in cases.



Syphilis Cases by Zip Code



Annual Comparisons



Risk Behaviors

- Multiple sex partners
- Bar or internet pick-ups
- Anonymous partners
- Sex for money or drugs
- Sex while using drugs or alcohol
- Lack of condom use


Multiple Partners

How do we target this population?

- Who are they?
 - 20-34 years of age
 - African American
 - Straight & men who have sex with men (MSM)
- Where/how do they meet?
 - Bars, restaurants, events
 - Internet, smart phone applications
 - Specific zip codes
 - Through friends, family, and acquaintances



Health Factors



Contributing Factors

Predisposing Factors

- Lack of awareness and knowledge of syphilis, preventive behaviors/methods, and available treatments
- Attitudes and beliefs
- Personal preferences

Reinforcing Factors

- Social support
- Social norms
- Economic reward
- Low perceived risk and susceptibility


Enabling Factors

- Lack of access to health care/testing
- No time for testing
- Income
- Environment

Stakeholders



Action Cycle



Actions

2013 Syphilis Awareness Campaign

- Awareness and knowledge
- Outreach testing
- Strengthen community partnerships
- Policy change

